

Package ‘imprProbEst’

February 14, 2012

Type Package

Title Minimum distance estimation in an imprecise probability model

Version 1.0.1

Date 2008-10-23

Author Robert Hable

Maintainer Robert Hable <Robert.Hable@uni-bayreuth.de>

Description A minimum distance estimator is calculated for an imprecise probability model. The imprecise probability model consists of upper coherent previsions whose credal sets are given by a finite number of constraints on the expectations. The parameter set is finite. The estimator chooses that parameter such that the empirical measure lies next to the corresponding credal set with respect to the total variation norm.

Depends R(>= 2.7.0), inline, lpSolve

LazyLoad yes

License LGPL-3

Repository CRAN

Date/Publication 2010-05-07 16:10:30

R topics documented:

imprProbEst-package	2
ArgMinDist	2

Index [5](#)

imprProbEst-package *Minimum distance estimation in an imprecise probability model*

Description

A minimum distance estimator is calculated for an imprecise probability model. The imprecise probability model consists of upper coherent previsions whose credal sets are given by a finite number of constraints on the expectations. The parameter set is finite. The estimator chooses that parameter such that the empirical measure lies next to the corresponding credal set with respect to the total variation norm.

Details

Package: imprProbEst
Type: Package
Version: 1.0
Date: 2008-10-23
License: LGPL-3
LazyLoad: yes

```
library(imprProbEst
```

Note

R programming support was given by Matthias Kohl

Author(s)

Robert Hable

Maintainer: Robert Hable <Robert.Hable@uni-bayreuth.de>

References

Hable (2008) *Data-Based Decisions under Complex Uncertainty*, Ph.D. thesis, LMU Munich, *in preparation*

Walley, P. (1991) *Statistical reasoning with imprecise probabilities*. Chapman & Hall, London.

Description

The function calculates a minimum distance estimator for an imprecise probability model. The imprecise probability model consists of upper coherent previsions whose credal sets are given by finite numbers of constraints on the expectations. The parameter set is finite. The estimator chooses that parameter such that the empirical measure lies next to the corresponding credal set with respect to the total variation norm.

Usage

```
ArgMinDist(x, lbomega, ubomega, epsilon, ImpreciseModel)
```

Arguments

<code>x</code>	a matrix where each row corresponds to one observation
<code>lbomega</code>	a vector containing the lower bounds of the sample space
<code>ubomega</code>	a vector containing the upper bounds of the sample space
<code>epsilon</code>	a positive real number; step size of the discretization
<code>ImpreciseModel</code>	a list of upper coherent previsions; see 'Details'

Details

The matrix `x` contains independent identically distributed data. Each row corresponds to one observation. The sample space is assumed to be a hyperrectangle in

$$R^k$$

. The lower bounds of this hyperrectangle are given by `lbomega`; the upper bounds of this hyperrectangle are given by `ubomega`. Accordingly, `length(lbomega)`, `length(ubomega)` and `length(x[, 1])` are equal to `k`.

Smaller values of `epsilon` may lead to more accurate results but increase the calculation time. Too small values of `epsilon` may cause an error due to RAM limitations.

`ImpreciseModel` contains an imprecise model consisting of upper coherent previsions. `ImpreciseModel` is a list; each component of `ImpreciseModel` is again a list which corresponds to an upper coherent prevision. Each upper coherent prevision is given by a list containing a list of functions and a corresponding vector of upper previsions.

For example, the imprecise model `ImpreciseModel` may consist of three coherent upper previsions `ImpreciseModel <- list(CohUpPrev1, CohUpPrev2, CohUpPrev3)`. `CohUpPrev1` may be defined by `CohUpPrev1 <- list(ListOfFunctions, UpperPrevisions)`. Here, `ListOfFunctions` is a list of functions, e.g., `ListOfFunctions <- list(f1, f2, f3, f4)`. Every function has to accept a single numeric argument and to return a numeric vector of the same length; the infimum of every function has to be 0, the supremum has to be 1. `UpperPrevisions` is a vector which contains the values of the upper coherent prevision at the functions in `ListOfFunctions`. That is, e.g., `UpperPrevisions[2]` is the value of the upper prevision at the function `f2`. Accordingly, the number of elements of the list `ListOfFunctions` is equal to `length(UpperPrevisions)`.

The estimation is that coherent upper prevision whose credal set has minimal total variation distance to the empirical measure generated by the observations `x`. Confer Hable (2008) for the definition of this minimum distance estimator; confer Walley (1991) and Hable (2008) for the theory of imprecise probabilities based on coherent upper previsions or coherent lower previsions.

Value

ArgMinDist returns a list, e.g. results, containing three components

results[[1]] the estimation; that is, the number of the minimizing coherent upper prevision in ImpreciseModel

results[[2]] the total variation distance of the minimizing coherent upper prevision

results[[3]] the number of linear programmes which had to be solved

Note

R programming support was given by Matthias Kohl

Author(s)

Robert Hable

References

Hable, R. (2008) *Data-Based Decisions under Complex Uncertainty*. Ph.D. thesis, LMU Munich, in preparation.

Walley, P. (1991) *Statistical reasoning with imprecise probabilities*. Chapman & Hall, London.

Examples

```
f1 <- function(v){ ifelse( abs(v-1)<1e-10 ,1,0) }
f2 <- function(v){ ifelse( abs(v-2)<1e-10 ,1,0) }
f3 <- function(v){ ifelse( abs(v-3)<1e-10 ,1,0) }
f4 <- function(v){ 1-ifelse( abs(v-3)<1e-10 ,1,0) }

x <- matrix(c(1,2,3,4),nrow=1)

UpperPrevisions1 <- c(1/4-0.03,1/4-0.03,1/4+0.01,1)
ListOfFunctions1 <- list(f1,f2,f3,f4)
CohUpPrev1 <- list(ListOfFunctions1,UpperPrevisions1)

UpperPrevisions2 <- c(1/4-0.04,1/4+0.04,1/4-0.01)
ListOfFunctions2 <- list(f1,f2,f3)
CohUpPrev2 <- list(ListOfFunctions2,UpperPrevisions2)

ImpreciseModel <- list(CohUpPrev1,CohUpPrev2)

lbomega <- 1
ubomega <- 4
epsilon <- 0.01

ArgMinDist(x,lbomega,ubomega,epsilon,ImpreciseModel)
```

Index

*Topic **nonparametric**

ArgMinDist, [2](#)

*Topic **package**

imprProbEst-package, [2](#)

ArgMinDist, [2](#)

imprProbEst (imprProbEst-package), [2](#)

imprProbEst-package, [2](#)